

Objectives: The impact of severe (>70%) contralateral carotid stenosis or occlusion (SCSO) on outcomes following carotid endarterectomy (CEA) is controversial. This study evaluated the effect of SCSO on outcomes following CEA.

Methods: All patients undergoing carotid endarterectomy at a single institution were retrospectively stratified according to the presence of SCSO. The peri-operative outcomes of CEA in the presence of SCSO were analyzed using univariate and multivariable methods.

Results: There were 2945 CEA performed from 1/89 to 1/06, including 736 (25%) patients with SCSO. Patients identified with SCSO had a higher rate of prior contralateral CEA (44% vs 21%; $P < .01$), positive intra-operative EEG changes (31% vs 17%; $P < .01$), and use of an intra-operative shunt (40% vs 28%; $P < .01$). Univariate analysis identified SCSO as a risk factor for any stroke (2.8% vs 1.5%; $P = .02$), death (2.2% vs 1.1%; $P = .03$), and any stroke/death (4.3% vs 2.4%; $P < .01$), but not ipsilateral stroke (1.6% vs 1.2%; $P = .4$). Multivariable regression demonstrated SCSO as an independent predictor of any stroke (OR1.8 [95% CI: 1.3-3.3]; $P = .05$) and any stroke/death (OR1.7 [95% CI: 1.1-2.7]; $P = .02$), without increasing risk of ipsilateral stroke (OR1.3 [95% CI: 0.6-2.7]; $P = .2$). The presence of SCSO was also associated with a higher risk of late mortality (HR1.3 [95% CI: 1.1-1.4]; $P < .01$).

Conclusions: Although the presence of SCSO is a risk factor for any stroke/death with CEA, it does not increase the risk of ipsilateral stroke. These data suggest that increased attention to peri-operative medical and hemodynamic management should be especially considered in this patient cohort as the observed strokes do not occur in the territory at risk from the surgical procedure.

Author Disclosures: D. C. Brewster: Nothing to disclose; R. P. Cambria: Nothing to disclose; M. F. Conrad: Nothing to disclose; J. Kang: Nothing to disclose; C. J. Kwolek: Nothing to disclose; G. M. LaMuraglia: Nothing to disclose; S. Mukhopadhyay: Nothing to disclose; V. I. Patel: Nothing to disclose.

PVSS8.

A Longitudinal Analysis of the Impact of Cumulative Intravenous Contrast Exposure on Renal Function in Patients With Occlusive and Aneurysmal Vascular Disease

Panos Kougiaris¹, Briana Lowery¹, Sherene Sharath¹, Peter H. Lin², Carlos Bechara¹, George Pisimisis¹, Taamee Pak², Neal R. Barshes¹. ¹Surgery, Michael E. DeBakey VAMC, Houston, Tex; ²Baylor College of Medicine, Houston, Tex

Objectives: Patients with occlusive or aneurysmal vascular disease are repeatedly exposed to intravenous (IV) contrast for diagnostic or therapeutic purposes. We sought to determine the long-term impact of cumulative iodinated IV contrast exposure (CIVCE) on renal function, the later defined with National Kidney Foundation (NKF) criteria.

Methods: We conducted a longitudinal analysis on consecutive patients without renal insufficiency at baseline (NFK stage I or II) who underwent interventions for

arterial occlusive or aneurysmal disease. We collected detailed data on any IV iodinated contrast exposure (including diagnostic or therapeutic angiography, cardiac catheterization, IV pyelography, computed tomography (CT) with IV contrast, and CT angiography); medication exposure throughout the observation period; comorbidities; and demographics. Primary end point was the development of renal failure (RF) (defined as NFK stage 4 or 5). Analysis was performed using a Cox Regression frailty model with clustering at the patient level.

Results: Patients ($n = 1298$) had a mean follow-up of 5.8 (range, 2.2-14) years. In the multivariate model with RF as dependent variable and after adjusting for the statistically significant covariates of baseline renal function (HR, 0.94; $P < .001$), age (HR, 0.42; $P < .001$), diabetes (HR, 2.2; $P = .013$), and congestive heart failure (HR, 3.2; $P < .001$), CIVCE remained an independent predictor of RF development (HR, 1.15; $P < .001$). In the multivariate survival analysis model and after adjusting for the statistically significant covariates of the newly developed RF (HR, 5.1; $P < .001$), perioperative myocardial infarction (HR, 5.2; $P = .004$), age (HR, 1.5; $P = .01$), total number of open operations (HR, 2.1; $P < .001$) and serum albumin (HR, 0.4; $P < .001$), CIVCE was an independent predictor of mortality (HR, 1.14; $P < .001$).

Conclusions: Cumulative IV iodinated contrast exposure is an independent predictor of renal failure and mortality in patients with occlusive and aneurysmal vascular disease.

Author Disclosures: N. R. Barshes: Nothing to disclose; C. Bechara: Nothing to disclose; P. Kougiaris: Nothing to disclose; P. H. Lin: Nothing to disclose; B. Lowery: Nothing to disclose; T. Pak: Nothing to disclose; G. Pisimisis: Nothing to disclose; S. Sharath: Nothing to disclose.

PVSS9.

Endothelial Function and Geometric Remodeling of Arterialized Vein Grafts Are Asynchronous

Warren J. Gasper, Hugh Alley, Maria Aleshin, Joseph Rapp, Marlene Grenon, Michael S. Conte, Christopher D. Owens. Vascular Surgery, University of California San Francisco, San Francisco, Calif

Objectives: Lower extremity vein bypass grafts (LEVBG) undergo geometric remodeling within the first month of surgery which directly correlates with subsequent patency. Veins are less vasoreactive than arteries and the extent of biochemical adaptation by vein grafts in the arterial environment is unknown. Flow-mediated vasodilation (FMD) measures endothelial-dependent vasodilation in response to hyperemia. We hypothesized that geometric remodeling of venous conduits would be correlated with endothelial function after LEVBG.

Methods: Patients undergoing LEVBG were prospectively enrolled. Brachial artery FMD was performed at baseline and vein graft FMD studies were performed at intervals following LEVBG.

Results: A total of 57 patients, median age 64 years (IQR, 61-71) with median follow-up 17 months (IQR, 10-32), were included. Most patients (91%) were male, 75% were white, 49% had diabetes, 81% were on a statin and the